

# SeaChange KFF™

Soil Balancing

Cropping

Pasture & Animal

Foliar & Fertigation

A blend of ocean essentials enhanced with fulvic acid and growth promotants.

# NATS NAN-THAN SOLD FORM Sea Change KFF<sup>TM</sup> Coase resembles + lable acid - No. St. St. Social grant) presents - In the Assess of the Social Social

# Description

A unique combination of kelp, fish and fulvic acid in a proven formula which is further boosted by the addition of plant growth promotants.

## Benefits

- Enhanced root growth.
- Natural chelating capacity.
- Increased flowering.
- Higher fruit to flower ratio.
- A rescue remedy in stress situations.
- Bio-stimulation for microbes & earthworms.

# **Packaging:**

- 1 L
- 5 L
- 20 I
- 200 L
- \_ 1000 L





### **APPLICATION RATES**

### **Foliar Spray:**

### **Vegetables**

1 L per 100 L water up to 5 L/ha equivalent Apply weekly or as required

### **Orchards & Vineyards**

1 L per 100 L water up to 7 L/ha equivalent Apply every 2 – 4 weeks or as required

### **Ornamentals & Turf**

1 L per 100 L water up to 5 L/ha equivalent Apply every 2 – 4 weeks or as required

### **Broadacre Crops**

2 – 3 L/ha in 80 – 100 L water, depending on canopy closure, apply as required

### **Pasture**

2 - 3 L/ha in 80 - 100 L water, apply as required

### **Spot-Spraying**

50 mL per 10 L water, apply as required

### **Fertigation:**

### **Vegetables**

10 – 20 L/ha

Apply weekly or as required

### **Orchards, Vineyards, Ornamentals &**

Turf 10 - 20 L/ha

Apply every 2 – 4 weeks or as required

### **Home Garden Application:**

Dilute 45 mL per 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre Apply fortnightly or as required

### **Tree Tubestock or Vegetable Seedling Treatment:**

Soak roots in 1:150 dilution

### **STORAGE & HANDLING**

Store in a cool dry area out of direct sunlight. This

product has been tested to store safely down to 5°C.
Crystallisation or sedimentation may occur below 5°C.
Do not pre-mix or store in diluted form. Seal lid/cap immediately after use.

TYPICAL ANALYSIS W/V	
Total Nitrogen (N) (as organic)	2.82%
Total Phosphorus (P) (as water soluble)	0.36%
Total Potassium (K) (as organic) Sulfur (S)	1.09%
Calcium (Ca)	0.96%
Magnesium (Mg)	0.14%
SG	0.10%
pH	1.1
Conductivity	2 – 4.5
Appearance	30 – 50 mS/cm
	Brown liquid

### **INSTRUCTIONS**

Shake or stir well before use. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application. Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

### **IMPORTANT**

- Contains some larger particles. Screening is recommended prior to application through fine filters.
- Some separation may occur over time. Agitate well during application to ensure solution is homogenous.
- Avoid spraying close to harvest if produce is susceptible to staining.
- Note: Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.



Phone: 02 6382 2165 1300 811 681

www.yladlivingsoils.com.au | www.yladlivingcompost.com.au 186 Milgadara Rd, Young NSW